

INTRODUCTION

The present publication is a monograph of *Motandra*, a genus of three species restricted to continental tropical Africa. The study is based mainly on herbarium material but the author has had the opportunity to study *M. guineensis* in the field in Ivory Coast and in Côte d'Ivoire.

PART XI A REVISION OF MOTANDRA A. DC. (APOCYNACEAE)

A. De GANDOLFI (1844) described *Motandra* from a specimen having its name published which has already been mentioned earlier by Gmelin in 1627. The name *Motandra* comes from the Greek word *μότανδρος*, penile, referring to the penicillate hairs at the base of the stamens.

Several new species were described about 1900, most of which have turned out to be synonymous with *M. guineensis*, while others belong to other related genera *Radermachera* and *Ocotea*.

The two other species were described by DURAND & PLANCHON in 1855 (*M. guineensis* Durand and *M. pseudopeltata* Durand) and by DURAND & DE KRUIF in 1901 and 1916 respectively.

A. P. M. DE KRUIF

Department of Plant Taxonomy
Wageningen Agricultural University, The Netherlands

Received 17-X-1983

Date of publication 20-IV-1984

The genus *Motandra* consists of three species, all of which are widely distributed species. *M. guineensis* is found in primary forests, secondary forests or gallery forests from Guinea-Congo to Uganda and Kenya. *M. hildebrandtii* and *M. pseudopeltata* are restricted to the savannas and woodlands and secondary forests of West and Central Africa. The area of distribution of both of the latter species partly overlaps that of *M. guineensis*.

HABIT AND CRIMINALITY

All *Motandra* species are hermaphrodites. The fruiting habit was observed in the field. *M. guineensis* bears the fruit on a peduncle which is bent either to the right or to the left, so it is not upright.

SYNONYMY

Motandra ssp. *guineensis* is the only species which flowers during the rainy season and fruits mature in the dry season.

INTRODUCTION

The present publication is a monograph of *Motandra*, a genus of three species restricted to continental tropical Africa. The study is based mainly on herbarium material but the author has had the opportunity to study *M. guineensis* in the field in Ivory Coast and in Cameroun.

HISTORY OF THE GENUS

A. DE CANDOLLE (1844) described the genus *Motandra*, basing it on *M. guineensis* which had already been described in *Echites* by THONNING in 1827. The name *Motandra* comes from the Greek μοτόν or μοτός, penicillum, referring to the penicillate hairs at the apex of the anthers.

Several new species were described about 1900, some of which have turned out to be synonymous with *M. guineensis*, while others belong to the closely related genera *Bassea* and *Oncinotis*.

The two other species which are maintained here, *M. lujaei* De Wild. & T. Durand and *M. poecilophylla* Wernham, were published in 1901 and 1916 respectively.

GEOGRAPHICAL DISTRIBUTION

The genus *Motandra* is restricted to West and Central Africa. The most widely distributed species, *M. guineensis* inhabits mostly secondary deciduous forests or gallery forests from Guinea(-Conakry) to Uganda and south to Angola. *M. lujaei* and *M. poecilophylla* are restricted to the more humid rain forests and secondary forests of Western Central Africa. The area of distribution of both of the latter species partly overlaps that of *M. guineensis*.

HABIT AND GROWTH

All *Motandra* species are lianas or climbing shrubs. As the present author observed in the field, *M. guineensis* shows the architectural model of TROLL (HALLÉ & OLDEMAN 1970: 125; — & TOMLINSON 1978: 242). This species winds either to the right as to the left, or is not twining.

FLOWERING SEASONS

Motandra species flower towards the end of the dry and the beginning of the rainy season and fruits mature in the dry season.

RELATIONSHIP TO OTHER GENERA

Motandra belongs to the subtribe *Baisseinae* of the tribe *Ichnocarpeae* of the *Apocynoideae*. The present author is of the opinion that the subtribe *Baisseinae* is a well defined natural unit, but he is not yet sure about the delimitation of the *Ichnocarpeae*, which could certainly be placed in the *Apocynoideae*. He will come back to the subject in one of his forthcoming publications.

Baisseinae Pichon ex De Kruif, subtribus nova; Pichon 1950: 112 (French description only).

Lignosae scandentes vel suffrutescens repentes herbis perennis similissimae, inermes, omnibus partis latice albo suffultae. Petioli latere adaxiale glandulas gerentes. Corolla aestivatione dextrorsa. Antherae conum pistili capitem obtectantem formantes, pedem connectivis pro parte cohaerentes. Ovarium semi-inferum disco quinquelobato circumdatum. Fructus folliculis duabus basi connatis latere adaxiale dehiscentibus compositus. Semina anguste ellipsoidea, non rostrata, apice coma pilis simplicibus suffulta.

Lianas, climbing shrubs or rhizomatous creepers with white latex in all parts. Spines, tendrils and stipules absent. Glands present at the adaxial side of the petiole. Secondary veins curved towards the margin, anastomosing; domatia or pits usually present in the axils of some secondary veins. Inflorescences axillary or terminal – sometimes both at the same time. Calyx lobes imbricate, almost free, often with colleters at the base. Corolla tube inside with tufts of hairs alternating with the stamens, corolla lobes in bud overlapping to the right. Stamens: anthers connivent in a cone around the gynoecium, adaxially pubescent, polliniferous in the upper part only, sagittate at the base, acuminate at the apex, coherent with the pistil head by a retinacle (an orbicular or nearly oblong patch inside on the connective). Pistil: ovary hemi-inferior; carpels two, connate at the extreme base only, abruptly narrowed into the style, surrounded by a disk, which is adnate to the ovary at the base only, with 5 sometimes obscure lobes, alternating with the stamens; pistil head consisting of a cylindrical lower portion, to which the 5 anthers adhere, a slightly 5-winged central portion and a bilobed apex. Fruit composed of 2 follicles, connate at the base, adaxially dehiscent with a longitudinal slit. Seeds narrowly ellipsoid, more or less laterally compressed, not beaked, with an apical coma; endosperm in a thin layer completely surrounding the embryo.

Type genus: *Baissea* A. DC.

3 genera:

Baissea A. DC. (14 species in tropical Africa).

Motandra A. DC. (3 species in tropical Africa).

Oncinotis Benth. (7 species in tropical Africa and Madagascar).

GENUS DIAGNOSIS

Motandra A. DC. 1844: 423; Bentham & Hooker 1876: 715; Baillon 1889: 206; Schumann 1895: 164; Stapf 1902: 224–226; 1904: 613–614; Pichon 1950: 113.

Type species: *M. guineensis* (Thonning) A. DC.

Climbing shrubs or lianas; white latex present. Branches unarmed, slightly sulcate when dry. Branchlets terete. Stipules none. Leaves opposite or subopposite, those of a pair equal, without colleters in the axils; petiole often rounded beneath and channeled above when dry, otherwise terete, glandular along the adaxial side; blade ovate to obovate or narrowly so, acuminate at the apex, rounded or less often slightly cordate at the base, entire, sparsely pubescent to almost glabrous on both sides; midrib pubescent, glabrescent on both sides, impressed above and prominent beneath; on each side 5–8(–9) conspicuous secondary veins, which are curved towards the margin, anastomosing, impressed above and prominent beneath; with tufts of hairs (domatia) in the axils of some secondary veins; tertiary venation inconspicuous, reticulate. Inflorescence terminal, thyrsoid, many-flowered. Bracts, and – if present – bracteoles, sepal-like or leafy, inside glabrous. Flowers 5-merous, actinomorphic or with only sepals unequal. Sepals imbricate, connate up to 1 mm at the base, subacute to obtuse at the apex, entire, outside densely pubescent, inside glabrous, ciliate, with inside at the base 0–3 colleters per sepal, situated at the margins; colleters elliptic to ovate or narrowly so. Corolla: tube with 5 deep dents, which are appearing as incrasations inside, from about half the length to the base of the lobes; tube outside shortly pubescent, except the portion covered by the sepals, inside glabrous, except for tufts of stiff hairs, alternating with the stamens; lobes in bud contorted and overlapping to the right, obtuse to subacute at the apex, recurved, entire, slightly ciliate. Stamens included, connivent in a cone, inserted on a slightly thickened ring at the base of the corolla tube; filaments obscurely s-shaped, abaxially glabrous, adaxially pubescent, most densely so in the upper half; anthers introrse, narrowly triangular, basifixed, acuminate and with penicillate hairs at the apex, sagittate at the base and at both tips provided with a subglobose incrasation which is up to 0.1 mm in diameter; cells two, discrete, parallel, dehiscent throughout by a longitudinal slit; between the tails inside on the connective an orbicular patch, the retinacle, 0.1–0.2 mm in diameter, with long hairs along the margin, adhering to the pistil head. Pistil: ovary hemi-inferior; carpels two, connate at the extreme base only, ovoid, laterally compressed, entirely or only apically densely pubescent, abruptly narrowed into the style, surrounded by a disk, adnate to the ovary at the base only, which is composed of a ring and 5 oblong lobes, alternating with the stamens; style obconical to almost cylindrical, composed of two connate strands; pistil head glabrous, consisting of a slightly 5-winged upper and a cylindrical lower portion, to the latter of which the 5 retinacles adhere; pistil head at the apex with two appendices which are united at the base, glabrous. Placentas adaxial, ovuliferous at the abaxial side

only. *Fruits* pendulous, composed of two (sub-)equal, spreading follicles, conate at their extreme base, dehiscent along an adaxial line of dehiscence, almost fusiform, with longest diameter below the middle, at the apex acute and when dry often curled; exocarp woody, with several conspicuous or obscure longitudinal wings, except at the adaxial side; endocarp stiff, thinly parchmentaceous, smooth inside. *Seeds* attached to the adaxial side of the fruit, with at the apex a dense tuft of simple hairs (coma), which is directed towards the apex of the fruit; grain flat, elliptic to ovate, glabrous, rounded at both ends, thickened at the margin with the hilum at the base and with a raphe ascending to the apex; testa very thin, smooth or minutely rugose; endosperm fleshy, in a very thin layer, which completely surrounds the embryo; embryo straight; cotyledons ovate or broadly so, cordate or less often cuneate at the base, rounded or sometimes slightly acuminate at the apex, glabrous on both sides, with 5–7 secondary veins; rootlet terete, rounded at the apex, glabrous.

Distribution: 3 species in West and Central Africa.

KEY TO THE SPECIES

1. Inflorescence congested; bracts conspicuous, mostly larger than the sepals, $1-16 \times 0.2-10$ mm; corolla $6.3-12.2$ mm long, tube $4.2-10.2$ mm long, lobes $2-2.7 \times 1-1.5$ mm; follicles $4-6 \times$ as long as wide, $4.5-7.5 \times 0.6-1.8 \times 0.6-1.5$ cm, longitudinally winged, wings $1-2$ mm wide in the mature fruit; seeds up to 25 in each follicle. From Equatorial Guinea to Angola (Cabinda) **2. *M. lujaei***
- Inflorescence rather lax; bracts inconspicuous, sepal-like, $0.5-1.8 \times 0.2-1$ mm **2**
2. Corolla $1.4-3$ mm long, tube $1.5-2$ mm long, lobes $1-1.5 \times 0.7-1.2$ mm; follicles $12-20 \times$ as long as wide, $3-10 \times 0.3-0.6 \times 0.3-0.7$ cm, obscurely longitudinally winged; wings up to 1 mm wide in the mature fruit; seeds up to 8 in each follicle. From Southern Cameroun to Congo-Brazzaville) **3. *M. poecilophylla***
- Corolla $4.8-10.6$ mm long, tube $2.5-4$ mm long, lobes $2.3-7 \times 1-2.1$ mm; follicles $2-9.5 \times$ as long as wide, $4-18 \times 1-3.5 \times 1-3.5$ cm, conspicuously longitudinally winged; wings $0.5-5$ mm wide in the mature fruit; seeds numerous. West and Central Africa. **1. *M. guineensis***

SPECIES DESCRIPTIONS

1. Motandra guineensis (Thonning) A. DC. 1844: 423; Bentham 1876: 74, pl. 1185; Schumann 1895: 164; Hiern 1898: 672–673; Hua 1902: 480–481; Stapf 1902: 224–225; Irvine 1930: 295; Hutchinson & Dalziel 1931: 47; Robijns 1947: 79–80; Irvine 1961: 628; Huber 1963: 80.

Fig. 1; Phot. 1; Map 1

Basionym: *Echites guineensis* Thonning 1827: 149; 1828: 169; Junghans 1961: 323; Hepper 1976: 27.


Type: Ghana: Aquapim (= Akwapim), Thonning 262 (C, holotype; isotype: P-JU 7118). Homotypic synonyms: *Echites paniculata* Thonning ex A. DC. 1844: 423 (not of Poiret 1811: 536–537, nor of Roxburgh 1832: 17–18). *Baissea guineensis* (Thonning) Roberty 1953: 1428.

Heterotypic synonyms: *M. rostrata* Schumann 1900: 307; Stapf 1902: 226; Schumann 1903: 318–319; Stapf 1904: 613. Type: Nigeria: Lagos State: Ishagama, Schlechter 12312 (holotype not seen, destroyed in B; lectotype: K; other isotypes seen: BR, PRE, Z).

M. pyramidalis Stapf 1902: 225; Pichon 1950: 113, 141, pl. IX, fig. 47. Type: Angola: Cuanza Norte: Alto Queta and Ambaca road, Welwitsch 5964 (BM, lectotype; isotypes: BR, C, COI, G, K, LISU, LD, MO, P, PRE).

M. altissima Stapf 1906: 526–527. Type: Uganda: Bunjiako (= Bunjako), Dawe 260 (K, lectotype; photographs: BR, GB, GH, LISC, NY, PRE, WAG).

Climbing shrub or liana, 0.50–40 m high climbing over shrubs and in trees, 5–40 m long or more. Trunk 1–10 cm in diameter; bark brown, first smooth, later longitudinally fissured. Branches pale brown, with small orange-brown lenticels; branchlets densely dark brown-pubescent, slightly glabrescent, sometimes with small pale brown lenticels at the woody base. Leaves: petiole 3.5–10 (–13) mm long, rusty brown-pubescent, glabrescent, along the adaxial side with small spherical glands on 0.3–1.5 mm long pubescent stalks, of which 2 mostly larger, 0.8–1.5 mm long stalked, and forming a pair; blade (1.8–) 2–4 (–5) × as long as wide, 3.5–14 × 1.4–4.6 cm, membranaceous to papery, or thinly coriaceous, medium to dark green above, paler beneath, glossy, sometimes ciliate with long hairs; blade often in the first pair or the first two pairs of a branchlet, suborbicular, rounded or emarginate at the apex and smaller, sometimes those of these pairs very unequal; domatia consisting of a tuft of pale brown hairs, sometimes absent. Inflorescence rather lax, 2.5–15.5 × 1.5–7 (–9) cm. Peduncle and branches rusty brown-pubescent, glabrescent. Pedicels 1–5 (–8) mm long, densely rusty brown-pubescent, glabrescent. Bracts, and – if present – bracteoles, ovate to triangular or narrowly so, 0.8–1.8 × 0.2–0.6 mm, subacute at the apex, outside rusty brown-pubescent, glabrescent, inside at the base often with 1–3 colleters, elliptic to ovate or narrowly so. Flowers sweet-scented. Sepals rusty brown, triangular, 1–3.8 × as long as wide, 1.0–1.9 × 0.3–1.2 mm; colleters 0–2 per sepal. Corolla in the mature bud 4–9 × as long as the calyx, 4.8–10.6 mm long; tube white to greenish-white, obconical, of which the basal 1–2 mm urceolate, 1.8–3.2 × as long as the calyx, 2.5–4 mm long, 2.5–3 mm in diameter




at the apex, 0.9–2 mm at the base; inside with tufts of hairs at 0.9–1.9 mm from the base; lobes white to greenish-white, narrowly ovate to narrowly obovate, 0.9–2 × as long as the tube, 2.5–5.5 × as long as wide, 2.3–7 × 1–2.1 mm, obscurely veined, outside glabrous or puberulous, usually only at the base, inside glabrous. *Stamens* inserted at 0.2–0.4 mm from the base of the tube; filaments 0.8–1.2 mm long; anthers 1.4–2.6 × as long as the filaments, 1.5–2.5 mm long, at the apex with 0.4–1 mm long hairs; tails 0.4–0.5 mm long; fertile portion 0.7–1 mm long. *Pistil* 2.5–3.1 mm long; carpels 0.8–1.2 × 0.5–0.9 × 0.2–0.3 mm, superior portion 0.4–0.7 mm long; disk: ring 0.1–0.3 mm high, lobes 0.3–0.7 × 0.3–0.5 mm, glabrous or only at the apex pubescent, sometimes with colleters, resembling those of the sepals, between the lobes; style 0.2–0.7



PHOT. 1. *Motandra guineensis* (Thonning) A. DC. (De Kruif 685, phot. LEEUWENBERG).

FIG. 1. *Motandra guineensis* (Thonning) A. DC.: 1. flowering branch, $\frac{2}{3} \times$; 2. part of branch, $\frac{2}{3} \times$; 3. leaf base with petiole from above, $4 \times$; 4. domatium, $4 \times$; 5. flower, $4 \times$; 6. opened flower showing 2 stamens and the pistil; the anther of the right stamen detached from the clavuncula; part of corolla, calyx, and 3 stamens removed, $20 \times$; 7. stamen, adaxial side, $6 \times$; 8. stamen, abaxial side, $6 \times$; 9. transverse section of ovary, $10 \times$; 10. pistil with longitudinal section of ovary, $10 \times$; 11. fruit, $\frac{2}{3} \times$; 12. immature fruit, $2 \times$; 13. seed, $\frac{2}{3} \times$; 14. embryo, $2 \times$. (1. Binuyo FHI 41203; 2. De Kruif 665; 3. De Kruif 685; 4, 11. De Kruif 722; 5–10. Leeuwenberg 11974; 12–14. Gossweiler 5223B).

mm long, glabrous; pistel head: upper portion $0.3-0.5 \times 0.3$ mm, lower portion $0.2-0.5 \times 0.3$ mm; appendices $0.5-1$ mm long, free portion $0.3-0.7$ mm long. Each placenta with 25–40 ovules. *Fruits* often with many lenticels, often subtended by the persistent calyx and bracts; follicles shaping an angle of about $130-180^\circ$, $2-9.5 \times$ as long as wide, $4-18 \times 1-3.5 \times 1-3.5$ cm; exocarp 1–2 mm thick, longitudinally winged; wings undulate, in the mature fruit 0.5–5 mm wide; exocarp at the outer surface with a 2–4 mm thick, deciduous, rusty brown tomentum, which may hide the wings. *Seeds* numerous; coma 29–78 mm long; grain $11-18.5 \times 5-8.5$ mm; testa smooth or minutely rugose; cotyledons broadly ovate, $4.9-9.5 \times 3-8$ mm; rootlet 2.5–3.5 mm long.



MAP 1. *Motandra guineensis* (Thonning) A. DC.

Distribution: West and Central Africa from Guinea (-Conakry) to Uganda and south to Angola.

Ecology: Mostly light or secondary deciduous forests or gallery forests and in secondary regrowth, on sand, clay and rocky outcrops; alt. 0–1200 m.

A selection of the about 350 specimens examined:

GUINEA (-Conakry): Komé (fl. May) Paroisse 164 (P); Macenta (fl. Mar.) Jacques-Félix 1549 (P); Kaoulendougou, near Nzo (fl. Mar.) Chevalier 21073 (P).

MALI: Kaba (Kangaba) (fl. May) Chevalier 13133 & 13242 (P).

SIERRA LEONE: Konno Town (fl.) Burbridge 481 (K); near Yifin (fl. Mar.) Morton & Gledhill SL 1150 (K, MO, WAG); Bunumbu (fl. Apr.) Deighton 3949 (K).

LIBERIA: Cess R., near Baitown (fr. Aug.) Baldwin 9013 (K).

IVORY COAST: Odienné (fr. Oct.) adam 27087 (MO); km 27 road Tai-Guiglo (imm. fr. July) Garnier PG/UB 32 (K); 17 km S of Daloa (fl. Mar.) Leeuwenberg 3058 (BR, C, COI, E, FHO, FI, GOET, K, L, LD, M, MO, P, PRE, U, UC, WAG, Z); km 13 road Divo-Tiassalé (Oct.) De Kruif 665 (UCI, WAG); Nzi-Noua, km 20 road Ndouci-Yamoussoukro (fl. Apr.) De Kruif 23 (UCI, WAG);

Bouroukro, on road Tanda-Koun-Fao (Oct.) *De Kruif* 680 (UCI, WAG); 8 km S. of Aniassué, along road to Akoupé (imm. fr. Oct.) *De Kruif* 685 (UCI, WAG).

GHANA: Tain Tributaries Block II For. Res. (fr. Dec.) *Adams* 2785 (K); Dedeswa (fl., FEb.) *Chipp* 145 (BM, FHO, K, Z); Bobira For. Res. (fl. Feb.) *Leeuwenberg* 11974 (WAG); Akin (= Achim) (fl.) *Dudgeon* 4 (P); Aquapim (= Akwapin) (fl.) *Thonning* 262 (C, P.JU, type); Shiare (fl. Apr.) *Hall* 1375 (K).

TOGO: Missahohé (fr. Nov.) *Mildbread* 7379 (HBG, K); Asrama, 30 km E. of Nuatja (fr. Sept.) *J. W. A. Jansen* 2658 (Wag).

BENIN REPUBLIC: Adja Ouéré (fl. Mar.) *Le Testu* 115 (BM, K, P); ibid. (fr. Dec.) *Le Testu* 272 (BM, P); Bassila Forest (fr. Dec.) *Poisson* 100 (K, P); Pobé (Sept.) *Adjano'houn* 102 (K, P).

NIGERIA: Oyo State: km 15 road Ibadan-Lagos (fl. Mar.) *Meikle* 1247 (B, BR, K, P); Gambari For. Res. (fl. Mar.) *Bernardi* 8725 (BR, G, K, S, WAG). Ogun State: 9 km N of Abeokuta (fl. May) *Roberty* 1593 (G, P). Lagos State: Ishagama (fl. Mar.) *Schlechter* 12312 (BR, K, PRE, Z, type of *M. rostrata*). Ondo State: Akure For. Res. (fl. Apr.) *Guile* 2595 (MO). Niger State: Abuja Idu, Kuje Distr. (fl. May) *Eimunjeze et al.* *FHI* 66455 (K, MO, WAG). Kaduna State: Sanga For. Res. (fr. Dec.) *Olorunfemi* *FHI* 55146 (K). Bendel State: Ubaja For. Res., Ishan Distr. (fl. Apr.) *Daramola* *FHI* 31263 (FHO, K). Anambra State: Onitsha (fl.) *Barter* 1636 (K). Cross River State: Iyamoyong For. Res., Obubra Distr. (fl. Apr.) *Binuyo* *FHI* 41203 (BR, FHO, K, WAG). Gongola State: Baissa (fl. Apr.) *Chapman* 5332 (K).

CAMEROUN: near Kumba (fr. Nov.) *Krukoff* 61 (MO); Ga, on road Mapou-Nganbé (fl. Aug.) *Letouzey* 7743 (P, YA); km 15 road Bertoua-Deng Deng (fl. Mar.) *Breteler* 1207 (A, BR, FI, G, K, LISC, M, P, UC, WAG, YA); Yaoundé (fl.) *Zenker & Staudt* 794 (BM, BR, COI, GOET, K, LE, M, NY, WU, S, Z); Nkolbisson (fl. Mar.) *Breteler* 2628 (A, BR, FI, K, M, P, UC, WAG, YA); ibid. (fl. Apr.) *De Wilde & De Wilde-Duyfjes* 2275 (BR, K, MO, P, PRE, WAG, YA, Z); ibid. (fl., imm. fr. May) *Leeuwenberg* 5731 (BR, K, P, WAG, YA); near Baréto (fl. Feb.) *Letouzey* 12011 (K, P, WAG, YA).

CENTRAL AFRICAN REPUBLIC: Bayanga (fl. Feb.) *Wraber* 26 (K); Boukoko (fl. Mar.) *Tisserant* 1111 (BM, BR, LISC, P, WAG); Bangui (fr. Oct.) *Breyne* 1593 (BR); Les Moroubas (fl. Mar.) *Tisserant* 1462 (BM, P, WAG); Yalinga (fl. Mar.) *Le Testu* 4608 (BM, BR, LISC, MO, P, WAG).

GABON: Tchibanga (fl. Oct.) *Le Testu* 1799 (BM, P).

ZAÏRE: Equateur: Boketa (fl. Mar.) *Evrard* 503 (BR); Likimi (imm. buds, Feb.) *Malchair* 194 (BR); Dundusana (fl. Sept.) *Mortehan* 444 (BR); Ifale R., Befale Terr. (fl. Feb.) *Evrard* 3485 (BR); Mondombe (fl., fr. Nov.) *Jespersen* 28 (BR); Djoa (imm. fr. May) *Evrard* (BR). Haut-Zaïre: Monga (fl. Feb.) *Lebrun* 2290 (BR, WAG); Kulu (fl., fr.) *Van den Brande* 136 (BR); Tukpo (fl. Apr.) *Gérard* 4517 (BR); Batande R., Doruma Terr. (fl. Apr.) *De Graer* 218 (BR); Bambesa (fl., fr. Mar.) *Gérard* 2288 (BR, P, WAG); Burumbu (fl. Nov.) *Bequaert* 1152 (BR, K); Weko, along road Ngazi-Bengamisa (fl., fr. Oct.) *Louis* 273 (BR, K, NY, P, UC); Yangambi (fr. July) *Louis* 10557 (BR, C, K, P); ibid. (fr. Feb.) *Louis* 13521 (B, BR, K, P); near Nala (fl. Mar.) *Seret* 790 (BR); Irumu (fl. Mar.) *Bequaert* 2937 (BR, K). Kivu: Semliki R., between Lesse and Vieux-Beni (fl. Mar.) *Bequaert* 3279 (BR); Mutongo (fl. May) *Gutzwiller* 2845 (BR, WAG); Bunyakiri (fl. Apr.) *Léonard* 4030 (BR, WAG). Bandundu: between Ipamu and Bandu (fl. Oct.) *Vanderijst* 12257 (BR). Bas-Zaïre: Matadi (fl. Feb.) *Dacremont* 344 (BR, P, WAG); Lovo, Songololo Terr. (imm. fr. Nov.) *Compère* 907 (BR, K).

SUDAN: Lotti Forest, Torit Distr. (fl. Mar.) *Jackson* 1281 (BM).

UGANDA: Bugoma Forest (buds, imm. fr.) *Dawe* 702 (K, paratype of *M. altissima*); Bunjako (fl.) *Dawe* 260 (K, lectotype of *M. altissima*).

BURUNDI: Usumbura (= Bujumbura) (fl. Feb.) *Léonard* 1602 (BR, WAG).

ANGOLA: Cabinda: near Buco Zau (fr. Aug.) *Gossweiler* 6597 (BM, COI, K, LISJC, LISU); ibid. (fr. July) *Gossweiler* 7284 (BR, COI, LISJC, LISU). Lunda: Dundo (fl.) *Gossweiler* 15 Oct. 1946 (B). Cuanza Norte: Alto Queto and Ambaca road (fl.) *Welwitsch* 5964 (BM, BR, C, COI, G, K, LD, LISU, MO, P, PRE, lectotype of *M. pyramidalis*); Golungo Alto (fr.) *Welwitsch* 1118 (BM, paratype of *M. pyramidalis*); Granja de S. Luiz (fl. Nov.) *Gossweiler* 5162 (BM, BR, COI, LISU); ibid. (fl.) *Gossweiler* 5223 (BM, COI, LISJC, LISU); ibid. (fr. Nov.) *Gossweiler* 5223B (BM, COI, LISJC, LISU); Fazenda (fr. Aug.) *Gossweiler* 5188 (BM, COI, LISJC, LISU).




FIG. 2. *Motandra lujaei* De Wild. & T. Durand: 1. flowering branch, $\frac{2}{3} \times$; 2. fruiting branch, $\frac{2}{3} \times$; 3. leaf base with petiole from above, $4 \times$; 4. domatia, $2 \times$; 5. flower, $6 \times$; 6. opened flower showing 1 stamen and the pistil; part of corolla, calyx and 4 stamens removed, $6 \times$; 7. stamen, adaxial side, $10 \times$; 8. seed, $\frac{2}{3} \times$. (1, 5-7. Gossweiler 6009; 2, 4, 8. N. Hallé & Le Thomas 161; 3. Luja 274).

Note: The lectotype of *M. pyramidalis*, Welwitsch 5964, is a composite number, consisting of 2 collections not far from each other, viz. Ambaca road and Alto Queta.


2. *Motandra lujaei* De Wild. & T. Durand 1901: 23–25; Staph 1902: 225–226; De Wildeman 1903: 17–18; Staph 1904: 613; De Wildeman 1906: 302, pl. 69; Pichon 1950: 113.

Fig. 2; Phot. 2; Map 2

Type: Zaïre: Bandundu: Lubue, *Luja* 274 (BR, lectotype; isotypes: K, P, S).

Climbing *shrub* or *liana*, 1–10 m high climbing over shrubs and in trees. *Branches* dark to medium brown when dry, with small lenticels; branchlets densely dark brown-pubescent, glabrescent, sometimes with small lenticels at the base. *Leaves*: petiole 3–10 mm long, rusty brown-pubescent, sometimes with long hairs, glabrescent, along the adaxial side with small spherical glands on 0.5–1.5 mm long stalks; blade (1.8–) 2–4 × as long as wide, 3.5–13 × 1.2 × 4.9 cm, papery; blade often in the first pair or the first two pairs of a branchlet suborbicular, rounded or emarginate at apex and smaller, sometimes those of these pairs very unequal; domatia consisting of a tuft of pale brown hairs. *Inflorescence* congested, 1.5–5 (–7.5) × 1–2.5 cm. Peduncle and branches dark brown-pubescent, sometimes also with some long hairs, glabrescent. Pedicels 1–4 mm long, dark brown-pubescent. Bracts numerous, ovate or narrowly so, 1–16 × 0.2–10 mm, obtuse to acuminate at the apex, cuneate to cordate at the base, outside dark brown-pubescent; bracteoles – if present – sepal-like. *Sepals* dark brown, ovate to triangular, 1.5–2 × as long as wide, 2–2.3 × 1.2–1.5 mm; colleters 0–3 per sepal. *Corolla* in the mature bud 2.5–6 × as long as the calyx, 6.3–12.2 mm long; tube white, less often yellow or dark brown, cylindrical, or nearly so, wider near the base, 1.9–5.1 × as long as the calyx, 4.2–10.2 mm long, 1–2.2 mm in diameter; inside with tufts of hairs at 1.9–2.8 mm from the base; lobes white, less often yellow or dark-brown, ovate to triangular, 0.2–0.5 × as long as the tube, 1.6–2 × as long as wide, 2–2.7 × 1–1.5 mm, outside densely pubescent and there often with glabrous patches, inside slightly pubescent or glabrous. *Stamens* inserted at about 0.2 mm from the base of the tube; filaments 1.2–1.6 (–2) mm long; anthers 1.2–2 × as long as the filaments, 2–2.5 mm long, with 0.1–0.5 mm long hairs at the apex; tails 0.4–0.5 mm long; fertile portion 1–1.2 mm long. *Pistil* 3–3.7 mm long; carpels 1–1.3 × 0.5–1.2 × 0.3–0.4 mm, superior portion 0.7–1 mm long; disk: ring 0.3–0.4 mm high, lobes 0.6–0.8 × 0.4–0.5 mm, pubescent; style 0.7–1.2 mm long, glabrous or sometimes slightly pubescent; pistil head: upper portion 0.2–0.5 × 0.3 mm, lower portion 0.3–0.4 × 0.3 mm; appendices 0.8–1 mm long, free portion 0.5–0.7 mm long. Each placenta with 10–25 ovules. *Fruits* often with many lenticels, often subtended by the persistent calyx and bracts; follicles shaping an angle of about 180°, 4–6 × as long as wide, 4.5–7.5 × 0.6–1.8 × 0.6–1.5 cm; exocarp up to 1 mm thick, longitudinally winged; wings undulate, in the mature fruit 1–2 mm wide; exocarp at the outer surface with a 1–2 mm thick,

deciduous, dark brown tomentum, which may hide the wings. Seeds up to 25 in each follicle; coma 15–30 mm long; grain 12–15 × 4–8 mm; testa minutely rugose; cotyledons ovate, 8–9.5 × 5–6 mm; rootlet 3–4 mm long.



MAP 2. *Motandra lujaei* De Wild. & T. Durand

Distribution: Equatorial Guinea to Angola (Cabinda).

Ecology: Rain forests or secondary forests, also in open places; alt. 0–900 m.



PHOT. 2. *Motandra lujaei* De Wild. & T. Durand (Caballé 259, phot. CABALLÉ).

Specimens examined:

EQUATORIAL GUINEA: Rio Muni: sin. loc (fl.) *Tessmann* 945 (K).
GABON: Minvoul (fl.) *Cottes* 30 Aug. 1906 (P); Acourenzorc (fl. Mar.) *Le Testu* 9060 (BM, LISC, P); Bélinga (fl. Oct.) *Caballé* 259 (WAG); ibid. (fr. July) *N. Hallé* 4199 (P); ibid. (fr. July) *N. Hallé & Le Thomas* 161 (P); 10 km NE of Lalara (fl. Sept.) *Breteler & De Wilde* 411/78 (WAG); Makokou (fl. June) *Farron* 7541 (P); ibid. (fl. Oct.) *Hladik* 1561 (P); ibid (fl. Feb.) *Hladik* 2575 (WAG); Lastoursville (fl. Apr.) *Le Testu* 7165 (BM, G, P); ibid. (fl. Apr.) *Le Testu* 18 Apr. 1929 (BM); ibid. (fl. Nov.) *Le Testu* 5 Nov 1929 (BM, BR, LISC, MO, US); km 16 road Lastoursville-Moanda (fl., fr. Sept.) *Breteler & De Wilde* 759/78 (WAG)

CONGO(-BRAZZAVILLE): N'Gouma, along Komono-M'Bila road (fl. Jan.) *Bouquet* 2397 (P).

ANGOLA: Cabina: Punga Munga (fl. Dec.) *Gossweiler* 6009 (BM, COI, LISJC, LISU).

ZAÏRE: Bandundu: Ipamu (fl. Sept.) *Vanderijst* 10798 & 10799 (BR); ibid. (fl. Oct.) *Vanderijst* 11145 (BR); Lubue (fl., imm. fr. May) *Luja* 274 (BR, K, P, S, lectotype); ibid. (fl., imm. fr. June) *Luja* 287 (BR, paratype); Idofa (fl. Oct.) *Vanderijst* 12606 (BR).

3. *Motandra poecilophylla* Wernham 1916: 228.

Type: Cameroun: Bitye, Bates 649 (BM, holotype; isotype: MO).

Fig. 3; Map 3

Climbing shrub or liana, 1–5 m high climbing, up to at least 6 m long. Branches greyish-brown, not lenticellate; branchlets with a deciduous, rusty brown, short tomentum, turning sparsely pubescent with age. Leaves: petiole 5–8 mm long, with a deciduous, rusty brown, short tomentum; blade 1.5–3.5 (–4) × as long as wide, 3–11 × 1.9–3.7 cm, papery to thinly coriaceous when dry, pale to medium green above, paler and sometimes with whitish-green patches beneath, glossy; blade sometimes rounded or emarginate at the apex and smaller, usually so in the first two pairs of leaves of a branchlet; domatia consisting of a tuft of rusty brown hairs. Inflorescence rather lax, 7–27 × 2–7 cm. Peduncle and branches with a deciduous, rusty brown, short tomentum. Pedicels 2–4 mm long, rusty brown-tomentellous. Bracts, and – if present – bracteoles, ovate to triangular or broadly so, 0.5–0.9 × 0.4–1 mm, subacute at the apex, outside rusty brown-tomentellous, inside at the base often with 1–3 colleters, elliptic to ovate. Sepals rusty brown, ovate to triangular or rhombic, 0.7–1.4 × as long as wide, 0.7–1.1 × 0.6–1.1 mm; colleters 0–3 per sepal. Corolla in the mature bud 1.5–4 × as long as the calyx, 1.4–3 mm long; tube white, sometimes with a yellowish-white throat, slightly obconical of which the basal 1 mm urceolate, 1.6–2.6 × as long as the calyx, 1.5–2 mm long; 0.7–1 mm in diameter at the apex, 1.2–1.5 mm at the base; inside with tufts of hairs at 0.7–1.2 mm from the base; lobes white, broadly ovate to broadly triangular, 0.5–0.7 × as long as the tube, 0.9–1.5 × as long as wide, 1–1.5 × 0.7–1.2 mm, obscurely veined, outside pubescent, usually only at the base, inside glabrous. Stamens inserted at about 0.2 mm from the base of the tube; filaments 0.5–0.7 mm long; anthers 1.5–2.6 × as long as the filaments, 1.1–1.5 mm long, with 0.2–0.6 mm long hairs at the apex; tails 0.3–0.4 mm long; fertile portion 0.6–0.7 mm long. Pistil 2–2.8 mm long; carpels 0.6–1 × 0.3–0.5 × 0.2–0.3 mm, superior portion 0.4–0.7 mm long; disk: ring 0.1–0.2 mm high, lobes 0.2–0.5 × 0.1–0.5 mm,





FIG. 3. *Motandra poecilophylla* Wernham: 1. flowering branch, $\frac{2}{3} \times$; 2. fruiting branch, $\frac{2}{3} \times$; 3. leaf base with petiole from above, $2 \times$; 4. domatia, $2 \times$; 5. flower, $10 \times$; 6. opened flower showing 1 stamen and the pistil; part of corolla, calyx, and 4 stamens removed, $20 \times$; 7. stamen, adaxial side, $20 \times$; 8. seed, $\frac{2}{3} \times$. (1. Florence 1697; 2-8. Letouzey 8276)

glabrous or only at the apex pubescent; style 0.2–0.4 mm long, glabrous; pistil head: upper portion 0.2–0.5 × 0.3 mm, lower portion 0.2–0.4 × 0.3 mm; appendices 0.6–1.2 mm long, free portion 0.3–0.9 mm long. Each placenta with 5–8 ovules. Fruits often subtended by the persistent calyx; follicles shaping an angle of about 90–180°, 12–20 × as long as wide, 3–10 × 0.3–0.6 × 0.3–0.7 cm; exocarp very thin, obscurely longitudinally winged, at the outer surface with an up to 1 mm thick, deciduous, rusty brown tomentum. Seeds up to 8 in each follicle; coma 20–35 mm long; grain 10–14 × 2–4 mm; rootlet 3–4 mm long.



MAP 3. *Motandra poecilophylla* Wernham.

Distribution: From Southern Cameroun to Congo.

Ecology: Rain forests or secondary forests, also in open places; alt. 0–700 m.

Specimens examined:

CAMEROUN: 15 km S of Ebolowa (fl., fr. Feb.) *De Wilde & De Wilde-Duyfjes* 1988A (WAG), 1988B (BR, K, MO, P, WAG, YA); N'Koemvone, km 14 road Ebolowa-Ambam (fl. Oct.) *J. J. F. E. de Wilde* 7591A (WAG); Ekouk (fr.) *Mildbraed* 5693 (HBG); between Kondébilong and Méyila, 53 km SE. of Mbalmayo (fl. May) *Asonganyi* 80 (WAG, YA); near Oveng, 30 km WNW of Sangméléma (fl., fr. July) *Letouzey* 11433 (BR, K, P, WAG, YA); Sangméléma (buds, June) *Hédin* 1271 (P); ibid. (buds, Mar.) *Meijer* 15289 (WAG); Mézésé, 20 km ENE of Sangméléma (fl. Oct.) *Letouzey* 8163 (BR, P, YA); near Méyos-Méla, 30 km ENS of Djoum (fl., fr. Nov.) *Letouzey* 8276 (BR, K, P, WAG, YA); Bitye (fl. Dec.) *Bates* 649 (BM, MO, type); ibid. (fl., fr. Oct.) *Bates* 1476 (HBG); ibid. (fl. Nov.) *Bates* 1526 (HBG); ibid. (fl. July) *Hédin* 1232 (P); Mekas, 72 km SE of Akonolinga (fl. July) *Asonganyi* 241 (WAG, YA); near curve of the Dja R. (fl. May) *Mildbraed* 5414(HBG); along road Minton I-Alati (fl. Jan.) *Letouzey* 11754 (P, YA).

EQUATORIAL GUINEA: Rio Muni: Campo area (fl. Mar.) Tessmann 508 (K).

GABON: Bélinga (buds, Dec.), *N. Hallé* 3599 (P); Lastoursville (fl., imm., fr., Mar.) *Le Testu* 7061 (BR, LISC, MO, P, WAG); ibid. (fl., fr. Apr.) *Le Testu* 7170 (BR, HBG, LISC, P, WAG); ibid. (fl., fr. Feb.) *Le Testu* 7940 (BM, BR, LISC, P, WAG); 10 km S of Makokou (fl., imm. fr. Mar.) *Florence* 1697 (WAG); SE of Makokou (fl. Feb.) *N. Hallé* 1045 (P); Lindémé (fl. Apr.) *Le Testu* 8018 (BR, HBG, P, WAG); sin. loc. (fl. July) *N. Hallé & Le Thomas* 3 (P).

CONGO(-BRAZZAVILLE): along road Mossendjo-Gabon (fl., fr. May) *F. Hallé* 1837 (P, WAG).

NOMEN NUDUM

M. stapfiana Mildbr. 1922: 80, 89. = *M. poecilophylla* Wernham. Specimens cited: *Mildbraed* 5414 (HBG) and 5693 (HBG).

EXCLUDED SPECIES

M. erlangeri Schumann 1903: 318. = *Oncinotis inandensis* J. M. Wood & M. Evans (Apocynaceae).

M. glabrata Baillon 1888: 760. = *Oncinotis glabrata* (Baillon) Stapf ex Hiern (Apocynaceae).

M. viridiflora Schumann 1903: 319. = *Bassea viridiflora* (Schumann) De Kruif, **comb. nov.** (Apocynaceae).

M. welwitschiana Baillon 1888: 760. = *Oncinotis hirta* Oliver (Apocynaceae).

REFERENCES

- BAILLON, H. 1888. Bull. Soc. Linn. Paris **1**.
- BAILLON, H. 1889. Histoire des Plantes **10**. – Hachette, Paris.
- BENTHAM, G. 1876. In: HOOKER, J. D., Hooker's Icones Plantarum **12**. – Longman, Rees, Orme, Brown, Green & Longman, London.
- BENTHAM, G. & J. D. HOOKER 1876. Genera Plantarum **2**. – Reeve & Co., London.
- DE CANDOLLE, A. 1844. Prodromus systematis naturalis regni vegetabilis **8**. – Fortin, Masson & Soc., Paris.
- DE WILDEMAN, E. 1903. Notes sur quelques Apocynacées laticifères de la Flore du Congo. – Bruxelles.
- DE WILDEMAN, E. 1906. Études de systématique et de géographie botanique sur la flore du Bas et du Moyen-Congo. Ann. Mus. Congo, Bot., ser. 5, **1**.
- DE WILDEMAN, E. & T. DURAND 1901. In: DURAND, T. & E. DE WILDEMAN, Materiaux pour la flore du Congo X. Bull. Soc. Bot. Belg. **40**: 7–41.
- HALLÉ, F. & R. A. A. OLDEMAN 1970. Essai sur l'architecture et la dynamique de croissance des arbres tropicaux. – Masson & Co., Paris.
- HALLÉ, F., R. A. A. OLDEMAN & P. B. TOMLINSON 1978. Tropical Trees and Forests. – Springer, Berlin-Heidelberg-New York.
- HEPPER, F. N. 1976. The West African Herbaria of Isert & Thonning. – Bentham-Moxon Trust, Kew, Richmond.
- HIERN, W. P. 1898. Catalogue of the African Plants collected by Dr. Friedrich Welwitsch in 1853–61. Dicotyledons. **1**(3). – London.
- HUA, H. 1902. Description des deux fruits d'Apocynacées africaines. Bull. Mus. Hist. Nat. Paris **8**: 479–481.
- HUBER, H. 1963. Apocynaceae. In: HEPPER, F. N. (ed.), Flora of West Tropical Africa, ed. 2, **2**. – HMSO, London.
- HUTCHINSON, J. & J. M. DALZIEL 1931. Flora of West Tropical Africa, ed. 1, **2**. – HMSO, London.
- IRVINE, F. R. 1930. Plants of the Gold Coast. – Oxford Univ. Press, London.
- IRVINE, F. R. 1961. Woody plants of Ghana. – Oxford Univ. Press, London.
- JUNGHANS, J. 1961. Thonning's and Isert's Collections from 'Danish Guinea' (Ghana) in West Tropical Africa. Botanisk Tidsskrift **57**.
- MILDBRAED, J. 1922. Wissenschaftliche Ergebnisse der Zweiten Deutschen Zentral-Afrika-Expedition 1910–1911, **2**. – Klinkhardt & Biermann, Leipzig.

- PICHON, M. 1950. Classification des Apocynacées: XXV Echitoïdées et XXVIII Supplément aux Plumérioidées. *Mem. Mus. Nat. Hist. Nat.*, ser. B, Bot. **1** (1): 1–174.
- POIRET, J. L. M. 1811. In: LAMARK, M., *Encyclopédie Méthodique Botanique. Supplément 2* (2). – Pandoucke, Paris.
- ROBERTY, G. 1953. Notes sur la flore de l'Ouest-africain. *Bull. Inst. Afr. Noire* **15**: 1396–1451.
- ROBIJNS, W. 1947. Flore des Spermatophytes du Parc National Albert, II Sympétales. – Govt. Bruxelles.
- ROXBURGH, W. 1832. *Flora Indica*, ed. 2, **2**. – Thacker & Co., Calcutta.
- SCHUMANN, K. 1895. Apocynaceae. In: ENGLER, A. & K. PRANTL (eds.), *Die natürlichen Pflanzenfamilien* **4**(2): 109–189.
- SCHUMANN, K. 1900. In: SCHLECHTER, R., *Westafrikanische Kautschuk-Expedition*. – Verlag des Kol.-Wirt. Komitees, Berlin.
- SCHUMANN, K. 1903. Apocynaceae africanae. In: ENGLER, A., *Bot. Jahrb.* **33**: 316–321.
- STAPF, O. 1902. Apocynaceae. In: THISELTON-DYER, W. T., *Flora of Tropical Africa* **4**(1): 24–231. – Reeve & Co., London.
- STAPF, O. 1904. Addenda. In: THISELTON-DYER, W. T., *Flora of Tropical Africa* **4**(1): 588–627. – Reeve & Co., London.
- STAPF, O. 1906. Plantae Novae Daweanae in Uganda lectae. *Journ. Linn. Soc.* **37**: 495–532.
- THONNING, P. 1827. In: SCHUMACHER, H. C. F., *Beskrivelse af Guineiske planter som ere funde af danske Botanikere, isaer af Etatsraad Thonning*. – Kjøbenhavn.
- THONNING, P. 1828. In: SCHUMANN, H. C. F., (The same paper in) *Det Kongelige danske Videnskabernes Selskabs naturvidenskabelige og mathematiske Afhandlinger* **3**.
- WERNHAM, H. F. 1916. New gamopetalae from the South Cameroons. *Journ. Bot.* **54**: 226–231.

ACKNOWLEDGEMENTS

I would like to thank staff and collaborators of the Department of Plant Taxonomy of Wageningen Agricultural University for all their help in connection with this monograph, especially Dr. A. J. M. LEEUWENBERG for his kind supervision, Miss Y. F. TAN for her fine drawings, Miss Drs. F. J. H. VAN DILST and Dr. C. E. JARVIS for the correction of the text. I am also very grateful to the directors and keepers of the following herbaria for putting their material at my disposal: A, B, BM, BR, C, COI, E, FHO, FI, G, GB, GH, GOET, HBG, K, L, LD, LE, LISC, LISJC, LISU, M, MO, NY, P, PRE, S, U, UC, UCI, US, W, WAG, WU, YA, Z.

I received a grant from the Wageningen Agricultural University Fund to support my field work in Ivory Coast; the Netherlands Foundation for the Advancement of Tropical Research (WOTRO) provided a grant that enabled me to carry out field work in Cameroun.

INDEX OF EXSICCATAE

The number in parentheses corresponds with the species-number of the revision. Only numbered collections have been listed.

- Adam, J. G., 4015, 27087 (1).
- Adams, C. D., 2476, 2785 (1).
- Adebusuyi, J. K., FHI 40937 (1).
- Adjanohoun, E. J., 102, 1484B, 1866B (1).
- Aké Assi, L., 402B, 2507, 5014B (1).
- Andoh, J. E., 2212 (1).
- Asonganyi, J. N., 80, 241 (3).

- Bagshawe, A. G., 1500 (1).
- Baldwin, J. T., jr., 9013, 13462, 13503, 13615, 13701 (1).
- Bamps, P., 1918 (1).
- Bamps, P., S. Martins & M. Silva, 4226 (1).
- Barter, C., 1636, 3411 (1).
- Bates, G. L., 649 (3), 1192 (1), 1476, 1526 (3).
- Bequaert, J. C., Zaïre: 1152, 2668, 2738, 2937, 2940, 3279, 6612 (1).
- Bernardi, L., 8725 (1).
- Binuyo, A., FHI 41203, 45470 (1).
- Bouquet, A., 2397 (2).
- Brande, van den, see Van den Brande.
- Breteler, F. J., 1207, 2628 (1).
- Breteler, F. J. & J. J. F. E. de Wilde, 411/78, 759/78 (2).
- Breteler, F. J., J. J. F. E. de Wilde & A. J. M. Leeuwenberg, 2387 (1).
- Breyne, H., 1593 (1).
- Brown, J. W., 684 (1).
- Brunel, 4591 (1).
- Burbridge, K., 481 (1).

- Caballé, G., 259 (2).
- Cansdale, C. S., 3978 (1).
- Cavaco, A., 1227 (1).
- Chandler, P., 1431 (1).
- Chapman, J. D., 5332 (1).
- Chevalier, A., 13133, 13242, 13935, 14102, 15138, 15139, 16874, 16923, 17717, 19090, 21073, 23242, 23396, 23800 (1).
- Chipp, T. F., 145 (1).
- Claessens, J., 419 (1).
- Compère, P., 907 (1).

- Dacremont, A., 344 (1).
- Dalziel, J. M., 23 (1).
- Daramola, B. O., FHI 31262, FHI 46015 (1).
- Daramola, B. O. & Osanyinlusi, FHI 90122 (1).
- Darko, K. O., 905 (1).
- Dawe, M. T., 260, 702 (1).

- Dawkins, H. C., 401 (1).
- De Giorgi, S., 1754 (1).
- De Graer, P. A. M., 218 (1).
- Deighton, F. C., 3942, 3949 (1).
- De Kruif, A. P. M., 23, 665, 680, 683, 685, 714, 715, 722, 724, 735, 791, 929 (1).
- Demeuse, F., 429 (1).
- Devred, R., 1255, 4044 (1).
- Dewèvre, A., 423, 995 (1).
- De Wilde, J. J. F. E., 7591A (3).
- De Wilde, W. J. J. O. & B. E. E. de Wilde-Duyfjes, 1988A, 1988B (3), 2275 (1).
- Donis, C., 1534, 1741, 2035, 2063, 2387 (1).
- Dudgeon, G. C., 4 (1).
- Dümmer, R., 1270, 2792, 4193, 4326, 5569 (1).

- Eggeling, W. F., 665 (1).
- Eimunjeze, Ekwuno & Onijamowo, FHI 69934 (1).
- Eimunjeze, Macauley & Adebusuyi, FHI 66455 (1).
- Emwiogbon, J. A., FHI 45347, FHI 61605 (1).
- Enti, A. A., SP 638 (1).
- Evrard, C., 503, 3485, 3936 (partly, mixture), 4096 (1).
- Eijnatten, van, see Van Eijnatten.

- Farmer, L., 483, 487, 489, 504 (1).
- Farron, C., 7541 (2).
- FHI, 3118, 13613, 22472, 22529, 23262, 27465, 28838, 29296, 31263, 37651, 40937, 41203, 45347, 45470, 46015, 55146, 61605, 66455, 67115, 69934, 70523, 70685, 90122 (1).
- Florence, J., 1697 (3).
- Foster, E. W., 195 (1).

- Garnier, P., PG/UB 32, PG/UB 109, PG/UB 137 (1).
- Gérard, P., 555, 2288, 2548, 3225, 3796, 3977, 4517, 5072, 5518 (1).
- Germain, R., 8190 (1).
- Gilbert, G., 2183, 7865 (1).
- Giorgi, de, see De Giorgi.
- Goossens, V., 4066 (1).
- Gossweiler, J., 484, 1133, 4442, 4448, 4534, 4775, 4776, 4845, 4871, 5097, 5098, 5160, 5162, 5162B, 5187, 5188, 5190, 5223, 5223B (1), 6009 (2), 6162, 6497, 6597, 7284, 10169 (1).

- Graer, de, see De Graer.
 Guile, D. P. M., 2595 (1).
 Gutzwiller, R., 2845 (1).
- Hall, J. B., 1375 (1).
 Hall, T., FHI 27465 (1).
 Hallé, F., 1837 (3).
 Hallé, N., 1045, 3599 (3), 4199 (2).
 Hallé, N. & A. Le Thomas, 3 (3), 161 (2).
 Hédin, N. & A. Le Thomas, 3 (3), 161 (2).
 Hédin, L., 1232, 1271 (3).
 Hladik, A., 1561, 2575 (2).
- Irvine, F. R., 339, 535, 832, 1807 (1).
- Jackson, J. K., 1281 (1).
 Jacques-Félix, H., 1549 (1).
 Jansen, J. W. A., 2658 (1).
 Jespersen, 28 (1).
 Johnson, W. H., 630, 642 (1).
 Jones, A. P. D., FHI 3118 (1).
- Kadoko, E. J. 39 (1).
 Keay, R. W. J., FHI 37651 (1).
 Keay, R. W. J. & A. Foglie, FHI 22529 (1).
 Keay, R. W. J., & T. A. Russell, FHI 22472 (1).
 Kennedy, J. D., 188 (1).
 Kitson, A. E., 1069 (1).
 Kruif, de, see De Kruif.
 Krukoff, B. A., 61, 75, 82 (1).
- Latilo, M. G. & B. O. Daramola, FHI 28838 (1).
 Latilo, M. G. & Oguntayo, FHI 70523 (1).
 Lebrun, J., 2290, 2495, 5175 (1).
 Leeuwenberg, A. J. M., 3058, 5731, 11245, 11951, 11974, 12005, 12159 (1).
 Léonard, A., 1602, 3963, 4030 (1).
 Le Testu, G., 115, 272, 1799, 1842, 4608 (1), 7061 (3), 7165 (2), 7170, 7940, 8018 (3), 9060 (2).
 Letouzey, R., 1668, 1842, 2879, 4370, 7743 (1), 8163, 8276, 11433, 11754 (3), 12011, 12213 (1).
 Louis, J., 273, 980, 1086, 1361, 2332, 2650, 3252, 3803, 5720, 5955, 6093, 6241, 6680, 8304, 8406, 10557, 12605, 13521, 13619, 13740, 13842, 13843, 15773 (1).
 Lowe, T., 1701 (1).
 Luja, E. P., 172 (1), 274, 287 (2).
- Macgregor, W. D., 592 (1).
 Maitland, T. D., 927 (1).
 Malchair, L., 194 (1).
 Maudoux, E., 40 (1).
 Meer, van, see Van Meer.
 Michelson, A., 216 (1).
- Meijer, W., 15162 (1), 15289 (3).
 Meikle, R. D., 930, 1247 (1).
 Mildbraed, J., 5087 (1), 5414, 5693 (3), 7379 (1).
 Millen, H., 114, 145 (1).
 Millson, A., 16 (1).
 Mortehan, M. G., 444 (1).
 Morton, J. K., GC 8572 (1).
 Morton, J. K. & D. Gledhill, SL 1150, SL 1794 (1).
- Odewo, T. K., FHI 67115 (1).
 Olorunfemi, J., FHI 55146 (1).
 Olorunfemi, J. & Fagbemi, FHI 70685 (1).
 Onochie, C. F. A., FHI 13613 (1).
- Paroisse, G., 164 (1).
 Poisson, E., 100 (1).
- Reygaert, F., 1208 (1).
 Roberty, G., 1593, 13852, 13965, 14085, 14168 (1).
 Rudier, N., 29 (1).
- Schlechter, R., 12312 (1).
 Schnell, R., 819, 903, 1100 (1).
 Seret, F., 790 (1).
 Serv. For Nigeria, 479 (1).
 Sillans, 1611 (1).
 Smythe, C. W., 266 (1).
 Swarbrick, J. T., (TS) 2607 (1).
 Synnott, T. J. H., 725 (1).
- Talbot, P. A., 2062 (1).
 Tamajong, S., FHI 23262 (1).
 Tessman, G., 508 (3), 945 (2).
 Thomas, A. S., D174 (1).
 Thonning, P., 262 (1).
 Tisserant, C., 811, 1111, 1387, 1462, 1611 = Sil-lans 1611, 1902 (1).
 Toussaint, L., 222 (1).
 Troupin, G., 7290, 7701, 9236, 9263 (1).
- Ujor, E., FHI 29296 (1).
 Unwin, A. H., 13, 126 (1).
- Van den Brande, J., 136, 401 (1).
 Vanderijst, H., 10798, 10799, 11145 (2), 12257 (1), 12606 (2).
 Van Eijnatten, C. L. M., 1125, 1284, 1603, 1673, 1691 (1).
 Van Meer, P. P. C., 695, 1053, 1073, 1090 (1).
 Verger, P., 1247, 1419 (1).
 Vermoesen, F. M. C., 1861, 1945 (1).
 Vigne, C., 1073, 2888, 3729, 4393 (1).
 Villiers, J. F., 623 (1).

Vroumsia Tchinaye, 40 VT (1).

Wagemans, J., 1736 (1).

Welwitsch, F., 723, 1118, 5964 (1).

Wilde, de, see De Wilde.

Wraber, T., 26 (1).

Zenker, G. & A. Staudt, 794 (1).

REGISTER

Synonyms are in *italics*. Page number of principal entries in **bold face**; those of figures and photographs in *italics*.

Apocynaceae Juss. 16

Apocynoideae Endl. 2

Baissea A. DC. 1, 2

– *guineensis* (Thonning) Roberty 5

– *viridiflora* (Schumann) De Kruif 16

Basseinae Pichon ex De Kruif 2

Echites P. Br. 1

– *guineensis* Thonning 5

– *paniculata* Thonning ex A. DC. 5

Ichnocarpeae Benth. & Hook. f. (sic; as subtribus) 2

Motandra A. DC. 1, 2, **3**

– *altissima* Stapf 5, 9

– *erlangeri* Schumann 16

– *glabrata* Baillon 16

– *guineensis* (Thonning) A. DC. 1, 3, 4, **5**, 6, 7, 8

– *lujaei* De Wild. & T. Durand 1, 4, **10**, **11**, 12

– *poecilophylla* Wernham 1, 4, **13**, **14**, 15, 16

– *pyramidalis* Stapf 5, 9, 11

– *rostrata* Schumann 5, 9

– *stapfiana* Mildbr. 16

– *viridiflora* Schumann 16

– *welwitschiana* Baillon 16

Oncinotis Bent. 1, 2

– *glabrata* (Baillon) Stapf ex Hiern 16

– *hirta* Oliver 16

– *inandensis* J. M. Wood & M. Evans 16